

## Product datasheet for **TA500639**

### SLC2A6 Mouse Monoclonal Antibody [Clone ID: OTI7E3]


#### Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI7E3
Applications:	IF, IHC, WB
Recommend Dilution:	IHC 1:200, WB: 1:200
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Full-length protein expressed in 293T cell transfected with human SLC2A6 expression vector
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1.47 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	54.5 kDa
Gene Name:	solute carrier family 2 member 6
Database Link:	<a href="#">NP_060055</a> <a href="#">Entrez Gene 11182</a> <a href="#">Human</a>
Background:	Hexose transport into mammalian cells is catalyzed by a family of membrane proteins, including SLC2A6, that contain 12 transmembrane domains and a number of critical conserved residues.
Synonyms:	GLUT6; GLUT9; HSA011372
Protein Families:	Transmembrane




[View online »](#)


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY SLC2A6 ([RC204391], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SLC2A6. Positive lysates [LY402605] (100ug) and [LC402605] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-SLC2A6 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human) (1:200).




Western blot analysis of extracts (10ug) from a mouse cell line and 3 different mouse tissues by using anti-SLC2A6 monoclonal antibody (1:200).




Immunohistochemical staining of paraffin-embedded colon tissue within the normal limits using anti-SLC2A6mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA500639, Dilution 1:50)



Immunohistochemical staining of paraffin-embedded endometrium tissue within the normal limits using anti-SLC2A6mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA500639, Dilution 1:50)



Immunohistochemical staining of paraffin-embedded prostate tissue within the normal limits using anti-SLC2A6 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA500639, Dilution 1:50)



Anti-SLC2A6 mouse monoclonal antibody (TA500639) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY SLC2A6 ([RC204391]).