

Product datasheet for **TA500888**

PDE10A Mouse Monoclonal Antibody [Clone ID: OTI4A10]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI4A10
Applications:	FC, WB
Recommend Dilution:	WB:1:500; FC:1:50
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human PDE10A (NP_006652) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	88.4 kDa
Gene Name:	phosphodiesterase 10A
Database Link:	NP_006652 Entrez Gene 10846 Human
Background:	Phosphodiesterases (PDEs) are a family of related phosphohydrolyases that selectively catalyze the hydrolysis of 3'cyclic phosphate bonds in adenosine and/or guanine 3',5' cyclic monophosphate (cAMP and/or cGMP). They regulate the cellular levels, localization and duration of action of these second messengers by controlling the rate of their degradation. There are 11 subtypes of PDEs, named PDE1-11; PDE4, 7 and 8 selectively degrade cAMP, PDE5, 6 and 9 selectively degrade cGMP and PDE1, 2, 3, 10 and 11 degrade both cyclic nucleotides. PDEs are expressed ubiquitously, with each subtype having a specific tissue distribution. These enzymes are involved in many signal transduction pathways and their functions include vascular smooth muscle proliferation and contraction, cardiac contractility, platelet aggregation, hormone secretion, immune cell activation, and they are involved in learning and memory.




[View online »](#)

Synonyms: FLJ11894; FLJ25677; HSPDE10A
Protein Families: Druggable Genome
Protein Pathways: Progesterone-mediated oocyte maturation, Purine metabolism

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PDE10A ([RC211181], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PDE10A. Positive lysates [LY416496] (100ug) and [LC416496] (20ug) can be purchased separately from OriGene.



HEK293T cells transfected with either [RC211181] overexpressing plasmid (Red) or empty vector control plasmid (Blue) were penetrated and immunostained by anti-PDE10A antibody (TA500888), and then analyzed by flow cytometry.