

Product datasheet for **TA501871**

ALDH1L1 Mouse Monoclonal Antibody [Clone ID: OTI6A10]


Product data:

| | |
|-------------------------|---|
| Product Type: | Primary Antibodies |
| Clone Name: | OTI6A10 |
| Applications: | FC, IHC, WB |
| Recommend Dilution: | WB 1:500~2000, FLOW 1:100 |
| Reactivity: | Human, Dog |
| Host: | Mouse |
| Isotype: | IgG1 |
| Clonality: | Monoclonal |
| Immunogen: | Full length human recombinant protein of human ALDH1L1 (NP_036322) produced in HEK293T cell. |
| Formulation: | PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide. |
| Concentration: | 0.69 mg/ml |
| Purification: | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G) |
| Predicted Protein Size: | 98.6 kDa |
| Gene Name: | aldehyde dehydrogenase 1 family member L1 |
| Database Link: | NP_036322 Entrez Gene 100855730 Dog Entrez Gene 10840 Human |
| Background: | The protein encoded by this gene catalyzes the conversion of 10-formyltetrahydrofolate, NADP, and water to tetrahydrofolate, NADPH, and carbon dioxide. The encoded protein belongs to the aldehyde dehydrogenase family and is responsible for formate oxidation in vivo. Deficiencies in this gene can result in an accumulation of formate and subsequent methanol poisoning. [provided by RefSeq] |
| Synonyms: | 10-fTHF; 10-FTHFDH; FDH; FTHFD |
| Protein Families: | Druggable Genome |
| Protein Pathways: | One carbon pool by folate |




[View online »](#)


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ALDH1L1 ([RC213720], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ALDH1L1. Positive lysates [LY415919] (100ug) and [LC415919] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-ALDH1L1 monoclonal antibody.




Immunohistochemical staining of paraffin-embedded Human colon tissue within the normal limits using anti-ALDH1L1 mouse monoclonal antibody. (TA501871)




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-ALDH1L1 mouse monoclonal antibody. (TA501871)




Immunohistochemical staining of paraffin-embedded Human liver tissue within the normal limits using anti-ALDH1L1 mouse monoclonal antibody. (TA501871)




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-ALDH1L1 mouse monoclonal antibody. (TA501871)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human pancreas tissue using anti-ALDH1L1 mouse monoclonal antibody. (TA501871)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human prostate tissue using anti-ALDH1L1 mouse monoclonal antibody. (TA501871)




Immunohistochemical staining of paraffin-embedded Human bladder tissue within the normal limits using anti-ALDH1L1 mouse monoclonal antibody. (TA501871)



Immunohistochemical staining of paraffin-embedded Carcinoma of Human bladder tissue using anti-ALDH1L1 mouse monoclonal antibody. (TA501871)



HEK293T cells transfected with either [RC213720] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-ALDH1L1 antibody (TA501871), and then analyzed by flow cytometry.



Flow cytometric Analysis of HeLa cells, using anti-ALDH1L1 antibody (TA501871), (Red), compared to a nonspecific negative control antibody ([TA50011]), (Blue).