

Product datasheet for TA502058

Annexin A3 (ANXA3) Mouse Monoclonal Antibody [Clone ID: OT11F11]


Product data:

Product Type:	Primary Antibodies
Clone Name:	OT11F11
Applications:	IHC, WB
Recommend Dilution:	WB 1:500~2000, IHC 1:150
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human ANXA3 (NP_005130) produced in HEK293T cell.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	36.2 kDa
Gene Name:	annexin A3
Database Link:	NP_005130 Entrez Gene 306 Human
Background:	This gene encodes a member of the annexin family. Members of this calcium-dependent phospholipid-binding protein family play a role in the regulation of cellular growth and in signal transduction pathways. This protein functions in the inhibition of phospholipase A2 and cleavage of inositol 1,2-cyclic phosphate to form inositol 1-phosphate. This protein may also play a role in anti-coagulation. [provided by RefSeq, Jul 2008]
Synonyms:	ANX3
Protein Families:	Stem cell - Pluripotency




[View online »](#)


Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ANXA3 ([RC201540], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ANXA3. Positive lysates [LY417500] (100ug) and [LC417500] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-ANXA3 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).



Western blot analysis of extracts (10ug) from 10 Human tissue by using anti-ANXA3 monoclonal antibody at 1:200 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: colon; 10: spleen).




Figure from citation: Western blot analysis of ANXA3 protein level by using anti-ANXA3 antibody in CD133+ HCC cell lines. HCC cells, including HepG2, SMMC-7721, and Hep3B cells. Dilution: 1:1000 [View Citation](#)






Figure from citation: Western Blot of ANXA3 protein level by using anti-ANXA3 antibody in human HCC tissues and nontumorous tissues. Dilution: 1:1000 [View Citation](#)



Immunohistochemical staining of paraffin-embedded Human lung tissue within the normal limits using anti-ANXA3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502058)



Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-ANXA3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502058)




Figure from citation: Representative images of immunohistochemical staining of ANXA3 expression in human liver tissues: (C) negative ANXA3 expression in non-tumorous liver parenchyma and (D) weak, (E) moderate, and (F) strong ANXA3 immunoreactivity in HCC tissues. Dilution: 1:500 [View Citation](#)