

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA502820

PNMT Mouse Monoclonal Antibody [Clone ID: OTI2E3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2E3

Applications: FC, IF, IHC, WB

Recommend Dilution: WB 1:2000, IHC 1:150, IF 1:100, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PNMT (NP_002677) produced in HEK293T

cell.

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.58 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 30.7 kDa

Gene Name: phenylethanolamine N-methyltransferase

Database Link: NP 002677 Entrez Gene 5409 Human

Background: The product of this gene catalyzes the last step of the catecholamine biosynthesis pathway,

which methylates norepinephrine to form epinephrine (adrenaline). The enzyme also has beta-carboline 2N-methyltransferase activity. This gene is thought to play a key step in

regulating epinephrine production. [provided by RefSeq]

Synonyms: PENT; PNMTase


Protein Families: Druggable Genome

Protein Pathways: Metabolic pathways, Tyrosine metabolism







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PNMT ([RC206586], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PNMT. Positive lysates [LY400946] (100ug) and [LC400946] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-PNMT mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502820)




HEK293T cells transfected with either [RC206586] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-PNMT antibody (TA502820), and then analyzed by flow cytometry.






Flow cytometric Analysis of Jurkat cells, using anti-PNMT antibody (TA502820), (Red), compared to a nonspecific negative control antibody, (Blue).




Flow cytometric Analysis of Hela cells, using anti-PNMT antibody (TA502820), (Red), compared to a nonspecific negative control antibody, (Blue).



Immunofluorescent staining of HeLa cells using anti-PNMT mouse monoclonal antibody (TA502820).





Anti-PNMT mouse monoclonal antibody (TA502820) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PNMT ([RC206586]).