

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA502998

# PDXK Mouse Monoclonal Antibody [Clone ID: OTI5H5]

## **Product data:**


Product Type:	Primary Antibodies
Clone Name:	OTI5H5
Applications:	FC, IF, IHC, WB
<b>Recommend Dilution:</b>	WB 1:2000, IHC 1:150, IF 1:100, FLOW 1:100
Reactivity:	Human
Host:	Mouse
lsotype:	lgG2a
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human PDXK(NP_03672) produced in HEK293 cell.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.82 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	34.9 kDa
Gene Name:	pyridoxal kinase
Database Link:	<u>NP_003672 Entrez Gene 8566 Human</u>
Background:	The protein encoded by this gene phosphorylates vitamin B6, a step required for the conversion of vitamin B6 to pyridoxal-5-phosphate, an important cofactor in intermediary metabolism. The encoded protein is cytoplasmic and probably acts as a homodimer. Alternatively spliced transcript variants have been described, but their biological validity has not been determined. [provided by RefSeq, Jul 2008]
Synonyms:	C21orf97; C21orf124; HEL-S-1a; PKH; PNK; PRED79
Protein Families:	Druggable Genome
Protein Pathways:	Metabolic pathways, Vitamin B6 metabolism




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US




### **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PDXK ([RC200975], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PDXK. Positive lysates [LY418499] (100ug) and [LC418499] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (10ug) from 7 different cell lines by using anti-PDXK monoclonal antibody (1:200).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US 


Equivalent amounts of cell lysates (10 ug per lane) of wild-type HEK293T cells (WT, Cat# LC810293T) and PDXK-Knockout HEK293T cells (KO, Cat# [LC841087]) were separated by SDS-PAGE and immunoblotted with anti-PDXK monoclonal antibody TA502998 (1:500). Then the blotted membrane was stripped and reprobed with anti-HSP90 antibody as a loading control.



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-PDXK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502998)

Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-PDXK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502998)


This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-PDXK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502998)


Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-PDXK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502998)

Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-PDXK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502998)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US 

Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-PDXK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502998)

Anti-PDXK mouse monoclonal antibody (TA502998) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PDXK ([RC200975]).



HEK293T cells transfected with either [RC200975] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-PDXK antibody (TA502998), and then analyzed by flow cytometry.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US