

Product datasheet for **TA504873**

ADH7 Mouse Monoclonal Antibody [Clone ID: OTI2H10]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI2H10
Applications:	FC, IHC, WB
Recommend Dilution:	WB 1:500~2000, IHC 1:150, FLOW 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human ADH7(NP_000664) produced in HEK293T cell.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	41.3 kDa
Gene Name:	alcohol dehydrogenase 7 (class IV), mu or sigma polypeptide
Database Link:	NP_000664 Entrez Gene 131 Human
Background:	This gene encodes class IV alcohol dehydrogenase 7 mu or sigma subunit, which is a member of the alcohol dehydrogenase family. Members of this family metabolize a wide variety of substrates, including ethanol, retinol, other aliphatic alcohols, hydroxysteroids, and lipid peroxidation products. The enzyme encoded by this gene is inefficient in ethanol oxidation, but is the most active as a retinol dehydrogenase; thus it may participate in the synthesis of retinoic acid, a hormone important for cellular differentiation. The expression of this gene is much more abundant in stomach than liver, thus differing from the other known gene family members. Alternative splicing results in multiple transcript variants. [provided by RefSeq]
Synonyms:	ADH4
Protein Families:	Druggable Genome




[View online »](#)

Protein Pathways: Drug metabolism - cytochrome P450, Fatty acid metabolism, Glycolysis / Gluconeogenesis, Metabolic pathways, Metabolism of xenobiotics by cytochrome P450, Retinol metabolism, Tyrosine metabolism


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ADH7 [RC224304], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ADH7. Positive lysates [LY424575] (100ug) and [LC424575] (20ug) can be purchased separately from OriGene.




HEK293T cells transfected with either [RC224304] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-ADH7 antibody (TA504873), and then analyzed by flow cytometry.



Immunohistochemical staining of paraffin-embedded Carcinoma of Human lung tissue using anti-ADH7 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA504873)



Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-ADH7 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA504873)



Western blot analysis of extracts (10ug) from 3 different cell lines by using anti-ADH7 monoclonal antibody (1:200).