

Product datasheet for TA505412


PTGES2 Mouse Monoclonal Antibody [Clone ID: OTI2C3]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI2C3
Applications:	IF, IHC, WB
Recommend Dilution:	WB 1:2000, IHC 1:150, IF 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human PTGES2(NP_079348) produced in HEK293T cell.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	41.8 kDa
Gene Name:	prostaglandin E synthase 2
Database Link:	NP_079348 Entrez Gene 80142 Human
Background:	The protein encoded by this gene is a membrane-associated prostaglandin E synthase, which catalyzes the conversion of prostaglandin H2 to prostaglandin E2. This protein also has been shown to activate the transcription regulated by a gamma-interferon-activated transcription element (GATE). Multiple transcript variants have been found for this gene. [provided by RefSeq]
Synonyms:	C9orf15; GBF-1; GBF1; mPGES-2; PGES2
Protein Pathways:	Arachidonic acid metabolism, Metabolic pathways




[View online »](#)

Product images:


HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PTGES2 ([RC221207], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PTGES2. Positive lysates [LY403045] (100ug) and [LC403045] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Human colon tissue within the normal limits using anti-PTGES2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA505412)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-PTGES2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA505412)




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-PTGES2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA505412)




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-PTGES2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA505412)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human thyroid tissue using anti-PTGES2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA505412)



Immunohistochemical staining of paraffin-embedded Human endometrium tissue within the normal limits using anti-PTGES2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA505412)



Immunohistochemical staining of paraffin-embedded Human bladder tissue within the normal limits using anti-PTGES2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA505412)



Anti-PTGES2 mouse monoclonal antibody (TA505412) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PTGES2 ([RC221207]).