

Product datasheet for TA505578


PGD Mouse Monoclonal Antibody [Clone ID: OTI2A5]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI2A5
Applications:	IF, IHC, WB
Recommend Dilution:	WB 1:2000, IHC 1:150, IF 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human PGD(NP_002622) produced in HEK293T cell.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	53 kDa
Gene Name:	phosphogluconate dehydrogenase
Database Link:	NP_002622 Entrez Gene 5226 Human
Background:	6-phosphogluconate dehydrogenase is the second dehydrogenase in the pentose phosphate shunt. Deficiency of this enzyme is generally asymptomatic, and the inheritance of this disorder is autosomal dominant. Hemolysis results from combined deficiency of 6-phosphogluconate dehydrogenase and 6-phosphogluconolactonase suggesting a synergism of the two enzymopathies. [provided by RefSeq]
Synonyms:	6PGD
Protein Pathways:	Glutathione metabolism, Metabolic pathways, Pentose phosphate pathway




[View online »](#)

Product images:


HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PGD ([RC201729], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PGD. Positive lysates [LY419202] (100ug) and [LC419202] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Human breast tissue within the normal limits using anti-PGD mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA505578)




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-PGD mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA505578)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-PGD mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA505578)




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-PGD mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA505578)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human thyroid tissue using anti-PGD mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA505578)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-PGD mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA505578)



Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-PGD mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA505578)



Immunohistochemical staining of paraffin-embedded Carcinoma of Human prostate tissue using anti-PGD mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA505578)



Anti-PGD mouse monoclonal antibody (TA505578) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PGD ([RC201729]).