

Product datasheet for TA505835

NRAS Mouse Monoclonal Antibody [Clone ID: OTI5G7]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI5G7
Applications:	IHC, WB
Recommend Dilution:	WB 1:2000, IHC 1:150
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human NRAS (NP_002515) produced in HEK293T cell.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	21 kDa
Gene Name:	NRAS proto-oncogene, GTPase
Database Link:	NP_002515 Entrez Gene 4893 Human
Background:	This is an N-ras oncogene encoding a membrane protein that shuttles between the Golgi apparatus and the plasma membrane. This shuttling is regulated through palmitoylation and depalmitoylation by the ZDHHC9-GOLGA7 complex. The encoded protein, which has intrinsic GTPase activity, is activated by a guanine nucleotide-exchange factor and inactivated by a GTPase activating protein. Mutations in this gene have been associated with somatic rectal cancer, follicular thyroid cancer, autoimmune lymphoproliferative syndrome, Noonan syndrome, and juvenile myelomonocytic leukemia. [provided by RefSeq, Jun 2011]
Synonyms:	ALPS4; CMNS; N-ras; NCMS; NRAS1; NS6
Protein Families:	Druggable Genome




[View online »](#)

Protein Pathways:

Acute myeloid leukemia, Axon guidance, B cell receptor signaling pathway, Bladder cancer, Chemokine signaling pathway, Chronic myeloid leukemia, Endometrial cancer, ErbB signaling pathway, Fc epsilon RI signaling pathway, Gap junction, Glioma, GnRH signaling pathway, Insulin signaling pathway, Long-term depression, Long-term potentiation, MAPK signaling pathway, Melanogenesis, Melanoma, Natural killer cell mediated cytotoxicity, Neurotrophin signaling pathway, Non-small cell lung cancer, Pathways in cancer, Prostate cancer, Regulation of actin cytoskeleton, Renal cell carcinoma, T cell receptor signaling pathway, Thyroid cancer, Tight junction, VEGF signaling pathway


Product images:


HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NRAS [RC202681], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NRAS. Positive lysates [LY400901] (100ug) and [LC400901] (20ug) can be purchased separately from OriGene.



Western blot analysis of anti-NRAS monoclonal antibody, TA505835. Incubation: 1:500, 1h.
 1: lysate of 293T transfected with HRAS plasmid, RC225202
 2: lysate of 293T transfected with NRAS plasmid, RC202681
 3: lysate of 293T transfected with KRAS plasmid, RC222697

HEK293T cells were transfected with the 3 different overexpression plasmids (1:HRAS, [RC225202]; 2: NRAS, [RC202681]; 3:KRAS, [RC222697]) for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-flag antibody ([TA180144], 1:1000) or anti-NRAS mouse monoclonal antibody. (TA505835, 1:500)



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-NRAS mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505835)