

Product datasheet for TA505872

MAPK4 Mouse Monoclonal Antibody [Clone ID: OTI2F4]


Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI2F4
Applications:	IF, IHC, WB
Recommend Dilution:	WB 1:4000, IHC 1:150, IF 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human MAPK4(NP_002738) produced in HEK293T cell.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	65.7 kDa
Gene Name:	mitogen-activated protein kinase 4
Database Link:	NP_002738 Entrez Gene 5596 Human
Background:	Mitogen-activated protein kinase 4 is a member of the mitogen-activated protein kinase family. Tyrosine kinase growth factor receptors activate mitogen-activated protein kinases which then translocate into the nucleus where it phosphorylates nuclear targets. [provided by RefSeq, Jul 2008]
Synonyms:	ERK-4; ERK4; p63-MAPK; p63MAPK; PRKM4
Protein Families:	Druggable Genome, Protein Kinase




[View online »](#)


Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MAPK4 ([RC207414], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MAPK4. Positive lysates [LY419131] (100ug) and [LC419131] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-MAPK4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505872)



Anti-MAPK4 mouse monoclonal antibody (TA505872) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY MAPK4 ([RC207414]).