

Product datasheet for **TA800688**


CD79A Mouse Monoclonal Antibody [Clone ID: OTI5E2]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI5E2
Applications:	IHC
Recommend Dilution:	IHC 1:150
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human CD79A (NP_001774) produced in HEK293T cell.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	21.8 kDa
Gene Name:	CD79a molecule
Database Link:	NP_001774 Entrez Gene 973 Human
Background:	The B lymphocyte antigen receptor is a multimeric complex that includes the antigen-specific component, surface immunoglobulin (Ig). Surface Ig non-covalently associates with two other proteins, Ig-alpha and Ig-beta, which are necessary for expression and function of the B-cell antigen receptor. This gene encodes the Ig-alpha protein of the B-cell antigen component. Alternatively spliced transcript variants encoding different isoforms have been described. [provided by RefSeq, Jul 2008]
Synonyms:	IGA; MB-1
Protein Families:	Druggable Genome, Transmembrane
Protein Pathways:	B cell receptor signaling pathway, Primary immunodeficiency




[View online »](#)

Product images:


Immunohistochemical staining of paraffin-embedded Carcinoma of Human lung tissue using anti-CD79A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA800688)



Immunohistochemical staining of paraffin-embedded Human lymph node tissue within the normal limits using anti-CD79A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA800688)



Immunohistochemical staining of paraffin-embedded Human lymphoma tissue using anti-CD79A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA800688)



Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-CD79A mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA800688)