

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA807052

Oxytocin neurophysin 1 (OXT) Mouse Monoclonal Antibody [Clone ID: OTI5G4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI5G4

Applications: WB

Recommend Dilution: WB 1:2000

Reactivity: Human Host: Mouse

Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 20-125 of human

OXT(NP_000906) produced in E.coli.

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 10.9 kDa

Gene Name: oxytocin/neurophysin I prepropeptide

Database Link: NP 000906 Entrez Gene 5020 Human

Background: This gene encodes a precursor protein that is processed to produce oxytocin and

neurophysin I. Oxytocin is a posterior pituitary hormone which is synthesized as an inactive precursor in the hypothalamus along with its carrier protein neurophysin I. Together with neurophysin, it is packaged into neurosecretory vesicles and transported axonally to the nerve endings in the neurohypophysis, where it is either stored or secreted into the bloodstream. The precursor seems to be activated while it is being transported along the axon to the posterior pituitary. This hormone contracts smooth muscle during parturition and lactation. It is also involved in cognition, tolerance, adaptation and complex sexual and maternal behaviour, as well as in the regulation of water excretion and cardiovascular

functions. [provided by RefSeq, Dec 2013]


Synonyms: OT; OT-NPI; OXT-NPI





Protein Families: Secreted Protein

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY OXT ([RC224226], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-OXT (1:2000). Positive lysates [LY424458] (100ug) and [LC424458] (20ug) can be purchased separately from OriGene.