

## Product datasheet for TA807178

### MNDA Mouse Monoclonal Antibody [Clone ID: OTI4H6]

#### Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI4H6
Applications:	IHC, WB
Recommend Dilution:	WB 1:2000, IHC 1:150
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 37-321 of human MNDA(NP_002423) produced in E.coli.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	45.7 kDa
Gene Name:	myeloid cell nuclear differentiation antigen
Database Link:	<a href="#">NP_002423</a> <a href="#">Entrez Gene 4332</a> <a href="#">Human</a>
Background:	The myeloid cell nuclear differentiation antigen (MNDA) is detected only in nuclei of cells of the granulocyte-monocyte lineage. A 200-amino acid region of human MNDA is strikingly similar to a region in the proteins encoded by a family of interferon-inducible mouse genes, designated Ifi-201, Ifi-202, and Ifi-203, that are not regulated in a cell- or tissue-specific fashion. The 1.8-kb MNDA mRNA, which contains an interferon-stimulated response element in the 5-prime untranslated region, was significantly upregulated in human monocytes exposed to interferon alpha. MNDA is located within 2,200 kb of FCER1A, APCS, CRP, and SPTA1. In its pattern of expression and/or regulation, MNDA resembles IFI16, suggesting that these genes participate in blood cell-specific responses to interferons. [provided by RefSeq, Jul 2008]
Synonyms:	PYHIN3




[View online »](#)

Protein Families: Transcription Factors


### Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MNDA ([RC207117], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MNDA. Positive lysates [LY400871] (100ug) and [LC400871] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human breast tissue using anti-MNDA mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA807178)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-MNDA mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA807178)




Immunohistochemical staining of paraffin-embedded Human liver tissue within the normal limits using anti-MNDA mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA807178)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human liver tissue using anti-MNDA mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA807178)




Immunohistochemical staining of paraffin-embedded Human Ovary tissue within the normal limits using anti-MNDA mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA807178)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-MNDA mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA807178)



Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-MNDA mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA807178)



Immunohistochemical staining of paraffin-embedded Human endometrium tissue within the normal limits using anti-MNDA mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA807178)



Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-MNDA mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA807178)