

Product datasheet for **TA809371**

XPB (ERCC3) Mouse Monoclonal Antibody [Clone ID: OTI6H8]


Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI6H8
Applications:	IHC, WB
Recommend Dilution:	WB 1:500, IHC 1:1000
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 191-436 of human ERCC3 (NP_000113) produced in E.coli.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	89.1 kDa
Gene Name:	ERCC excision repair 3, TFIIH core complex helicase subunit
Database Link:	NP_000113 Entrez Gene 2071 Human
Background:	ERCC3 is an ATP-dependent DNA helicase that functions in nucleotide excision repair and complements xeroderma pigmentosum group B mutations. It also is the 89 kDa subunit of basal transcription factor 2 (TFIIH) and thus functions in class II transcription. [provided by RefSeq, Jul 2008]
Synonyms:	BTF2; GTF2H; RAD25; TFIIH; TTD2; XPB
Protein Families:	Druggable Genome, Transcription Factors
Protein Pathways:	Nucleotide excision repair




[View online »](#)


Product images:




Western blot analysis of extracts (35ug) from 5 different cell lines by using anti-ERCC3 monoclonal antibody (1:500).




Immunohistochemical staining of paraffin-embedded Human colon tissue within the normal limits using anti-ERCC3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA809371) (1:1000)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-ERCC3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA809371) (1:1000)




Immunohistochemical staining of paraffin-embedded Human lung tissue within the normal limits using anti-ERCC3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA809371) (1:1000)




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-ERCC3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA809371) (1:1000)



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-ERCC3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA809371) (1:1000)



Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-ERCC3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA809371) (1:1000)



Immunohistochemical staining of paraffin-embedded Carcinoma of Human prostate tissue using anti-ERCC3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA809371) (1:1000)