

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA809596

### Pepsinogen II (PGC) Mouse Monoclonal Antibody [Clone ID: OTI5C9]

#### **Product data:**

| Product Type:              | Primary Antibodies  |
|----------------------------|---|
| Clone Name:                | OTI5C9  |
| Applications:              | LMNX  |
| <b>Recommend Dilution:</b> | 1:100 - 1:1000  |
| Reactivity:                | Human   |
| Host:                      | Mouse   |
| lsotype:                   | lgG1  |
| Clonality:                 | Monoclonal  |
| Immunogen:                 | Human recombinant protein fragment corresponding to amino acids 17-388 of human<br>PGII(NP_002621) produced in E.coli.  |
| Formulation:               | PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.  |
| Concentration:             | 1 mg/ml   |
| Purification:              | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography<br>(protein A/G)  |
| Predicted Protein Size:    | 40.5 kDa  |
| Gene Name:                 | progastricsin (pepsinogen C)  |
| Database Link:             | <u>NP_002621 Entrez Gene 5225 Human</u>   |
| Background:                | This gene encodes an aspartic proteinase that belongs to the peptidase family A1. The encoded protein is a digestive enzyme that is produced in the stomach and constitutes a major component of the gastric mucosa. This protein is also secreted into the serum. This protein is synthesized as an inactive zymogen that includes a highly basic prosegment. This enzyme is converted into its active mature form at low pH by sequential cleavage of the prosegment that is carried out by the enzyme itself. Polymorphisms in this gene are associated with susceptibility to gastric cancers. Serum levels of this enzyme are used as a biomarker for certain gastric diseases including Helicobacter pylori related gastritis. Alternate splicing results in multiple transcript variants. A pseudogene of this gene is found on chromosome 1. [provided by RefSeq, Oct 2009] |
| Synonyms:                  | PEPC; PGII  |




View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

**Protein Families:** 

Protease, Secreted Protein

## **Product images:**



PGII Luminex with 5C9 Capture (TA809596) and 1H5 Detection ([TA809141]) Antibodies. Substrate used: recombinant protein expressed in E.coli corresponding to amino acids 17-388 of human pepsinogen C (PG II).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US